

ON THE INTEGRAL KERNELS OF DERIVATIVES OF THE ORNSTEIN-UHLENBECK SEMIGROUP

JONAS TEUWEN

Abstract. This paper presents a closed-form expression for the integral kernels associated with the derivatives of the Ornstein-Uhlenbeck semigroup e^{tL} . Our approach is to expand the Mehler kernel into Hermite polynomials and applying the powers L^N of the Ornstein-Uhlenbeck operator to it, where we exploit the fact that the Hermite polynomials are eigenfunctions for L . As an application we give an alternative proof of the kernel estimates by [11], making all relevant quantities explicit.

1. Introduction

Much effort [2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13] has gone into developing the harmonic analysis of the *Ornstein-Uhlenbeck operator*

$$(1) \quad L := \frac{1}{2}\Delta - \langle x, \nabla \rangle.$$

On the space $L^2(\mathbf{R}^d, d\gamma)$, where γ is the Gaussian measure

$$(2) \quad d\gamma(x) := \pi^{-d/2} e^{-|x|^2} dx,$$

this operator can be viewed as the Gaussian counterpart of the Laplace operator Δ . Indeed, one has $L = -\nabla^* \nabla$, where ∇ is the usual gradient and the ∇^* is its adjoint in $L^2(\mathbf{R}^d, d\gamma)$. It is a classical fact that the semigroup operators e^{tL} , $t > 0$, are integral operators of the form

$$e^{tL}u(\cdot) = \int_{\mathbf{R}^d} M_t(\cdot, y)u(y) d\gamma(y),$$

where M_t is the so-called *Mehler kernel* [12]

$$(3) \quad M_t(x, y) = \frac{\exp\left(-\frac{|e^{-t}x - y|^2}{1 - e^{-2t}}\right)}{(1 - e^{-2t})^{d/2}} e^{|y|^2}$$

(see [13] for a representation of M_t which makes the symmetry in x and y explicit).

Developing a Hardy space theory for L is the subject of active current research [11, 6]. In this theory the derivatives $(d^k/dt^k)e^{tL} = L^N e^{tL}$ play an important role. The aim of the present paper is to derive closed form expressions for the integral kernels of these derivatives, that is, to determine explicitly the kernels M_t^N such that we have the identity

$$(4) \quad L^N e^{tL}u(\cdot) = \int_{\mathbf{R}^d} M_t^N(\cdot, y)u(y) d\gamma(y).$$

Direct application of the derivatives d^N/dt^N to the Mehler kernel yields expressions which become intractable even for small values of N . Our approach will be to

expand the Mehler kernel in terms of the L^2 -normalised Hermite polynomials and then to apply L^N to it, thus exploiting the fact that the Hermite polynomials are eigenfunctions for L .

As an application of our main result, which is proved in section 5 after developing some preliminary material in the sections 2-4, we shall give a direct proof for the kernel bounds of [11] in section 6.

2. Preliminaries

In this preliminary section we collect some standard properties of Hermite polynomials and their relationship with the Ornstein-Uhlenbeck operator. Most of this material is classical and can be found in [12, 14].

2.1. Hermite polynomials. The *Hermite polynomials* H_n , $n \geq 0$, are defined by Rodrigues's formula

$$(5) \quad H_n(x) := (-1)^n e^{x^2} \partial_x^n e^{-x^2}.$$

Their L^2 -normalizations,

$$h_n := \frac{H_n}{\sqrt{2^n n!}}$$

form an orthonormal basis for $L^2(\mathbf{R}, d\gamma)$. We shall use the fact that the generating function for the Hermite polynomials is given by

$$(6) \quad \sum_{n=0}^{\infty} \frac{H_n(x)}{n!} t^n = e^{2tx - t^2}.$$

The relationship with the Ornstein-Uhlenbeck operator is encoded in the eigenvalue identity $LH_n = -nH_n$, from which it follows that for all $t \geq 0$ we have $e^{tL}H_n = e^{-tn}H_n$. From this one quickly deduces that the Mehler kernel is given by

$$(7) \quad M_t(x, y) := \sum_{n=0}^{\infty} e^{-tn} h_n(x) h_n(y).$$

We will need two further identities for the Hermite polynomials which can be found, e.g., in [1, Chapter 18]: the integral representation

$$(8) \quad H_n(x) = \frac{(-2i)^n e^{x^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} \xi^n e^{2ix\xi} e^{-\xi^2} d\xi$$

and the “binomial” identity

$$(9) \quad H_n(x+y) = \sum_{k=0}^n \binom{n}{k} (2y)^{n-k} H_k(x).$$

2.2. Hermite polynomials in several variables. The Hermite polynomials on $\mathbf{R}^d$ are defined, for multiindices $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbf{N}^d$ by the tensor extensions

$$(10) \quad H_\alpha(x) := \prod_{i=1}^d h_{\alpha_i}(x_i),$$

for $x = (x_1, \dots, x_d)$ in $\mathbf{R}^d$. The normalized Hermite polynomials

$$(11) \quad h_\alpha := \frac{H_\alpha}{\sqrt{2^{|\alpha|} \alpha!}}, \quad \text{where } \alpha! = \alpha_1! \cdots \alpha_d!$$

form an orthonormal basis in $L^2(\mathbf{R}^d, d\gamma)$, and we have the eigenvalue identity

$$(12) \quad LH_\alpha = -|\alpha|H_\alpha, \quad \text{where } |\alpha| = \alpha_1 + \dots + \alpha_d.$$

If we consider the action of $L^N e^{tL}$ on a Hermite polynomial h_α , through the multinomial theorem applied to $|\alpha|^k$ we get (writing L_d for the operator L in dimension d and L_1 for the operator L in dimension 1)

$$(13) \quad \begin{aligned} L_d^N e^{tL} h_\alpha(x) &= |\alpha|^N e^{-t|\alpha|} h_{\alpha_1}(x_1) \cdot \dots \cdot h_{\alpha_d}(x_d) \\ &= \sum_{|n|=N} \binom{N}{n_1, n_2, \dots, n_d} \alpha_1^{n_1} \cdot \dots \cdot \alpha_d^{n_d} e^{-t\alpha_1} \cdot \dots \cdot e^{\alpha_d} h_{\alpha_1}(x_1) \cdot \dots \cdot h_{\alpha_d}(x_d) \\ &= \sum_{|n|=N} \binom{N}{n_1, n_2, \dots, n_d} L_1^{n_1} e^{tL_1} h_{\alpha_1}(x_1) \cdot \dots \cdot L_1^{n_d} e^{tL_1} h_{\alpha_d}(x_d). \end{aligned}$$

This implies that we can reduce the question of computing the d -dimensional version of the integral kernel to the one-dimension one.

3. A combinatorial lemma

From now on we concentrate on the Ornstein-Uhlenbeck operator L in one dimension, i.e., in $L^2(\mathbf{R}, d\gamma)$. We are going to follow the approach of [12]. Recalling the identity $Lh_n = -nh_n$, we will apply L^N to the generating function of the Hermite polynomials (6). A problem which immediately occurs is that Δ and $\langle x, \nabla \rangle$ do not commute, and because of this we cannot use a standard binomial formula to evaluate L^N . Instead, we note that

$$Lg = -t\partial_t g.$$

In particular this implies that

$$L^N g = (-1)^k D_N g,$$

where

$$(14) \quad D_N := \underbrace{t\partial_t \circ t\partial_t \circ \dots \circ t\partial_t}_{k \text{ times}} = (t\partial_t)^k.$$

The following lemma will be very useful.

Lemma 1. *We have*

$$(15) \quad D_N = \sum_{n=0}^N \left\{ \begin{matrix} N \\ n \end{matrix} \right\} t^n \partial_t^n,$$

where $\left\{ \begin{matrix} N \\ n \end{matrix} \right\}$ are the Stirling numbers of the second kind.

The Stirling numbers of the second kind are quite well-known combinatorial objects. For the sake of completeness we will state their definition and recall some relevant properties below. For more information we refer the reader to [15]. The related Stirling numbers of the first kind will not be needed here.

We begin by recalling the definition of *falling factorial*

$$(16) \quad (j)_n := j(j-1) \dots (j-n+1) = \frac{j!}{(j-n)!},$$

for non-negative integers $k \geq n$.

Definition 1. For non-negative integers $N \geq n$, the number Stirling number of the second kind $\left\{ \begin{smallmatrix} N \\ n \end{smallmatrix} \right\}$ is defined as the number of partitions of an N -set into n non-empty subsets.

These numbers satisfy the recursion identity

$$(17) \quad \left\{ \begin{smallmatrix} N \\ n \end{smallmatrix} \right\} = N \left\{ \begin{smallmatrix} N-1 \\ n \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} N-1 \\ n-1 \end{smallmatrix} \right\}.$$

For all non-negative integers j and k one has the generating function identity

$$(18) \quad j^N = \sum_{n=0}^N \left\{ \begin{smallmatrix} N \\ n \end{smallmatrix} \right\} (j)_n.$$

4. Weyl Polynomials

Before turning to the proof of lemma 1, let us already mention that it only depends on the commutator identity $[t, \partial_t] = -1$. This brings us to the observation that *Weyl polynomials* provide the natural habitat for our expressions. Roughly speaking, a Weyl polynomial is a polynomial in two non-commuting variables x and y which satisfy the commutator identity $[x, y] = -1$. This is made more precise in the following definition.

Definition 2. The Weyl algebra over a field $\mathbf{F}$ of characteristic zero is the ring $\mathbf{F}\langle x, y \rangle$ of all polynomials of the form $p(x, y) = \sum_{m=0}^M \sum_{n=0}^N c_{mn} x^m y^n$ with coefficients $c_{mn} \in \mathbf{F}$ in two noncommuting variables x and y which satisfy the commutator identity

$$[x, y] := xy - yx = -1.$$

We now have the following abstract version of lemma 1:

Lemma 2. In the Weyl algebra $\mathbf{F}\langle x, y \rangle$ we have the identity

$$(19) \quad (xy)^m = \sum_{i=1}^m \left\{ \begin{smallmatrix} m \\ i \end{smallmatrix} \right\} x^i y^i,$$

where $\left\{ \begin{smallmatrix} m \\ i \end{smallmatrix} \right\}$ are the Stirling numbers of the second kind.

As a preparation for the proof of lemma 2 we make a couple of easy computations. If we set $D := xy$, then

$$(20) \quad Dx^m = x^m D + mx^m,$$

$$(21) \quad Dy^m = y^m D - my^m.$$

This can be shown by induction on m . For instance, note that

$$Dx^m = x(D+1)x^{m-1} = xDx^{m-1} + x^m.$$

If we take this a bit further and have $p \in \mathbf{F}[D]$, then

$$(22) \quad p(D)x^m = x^m p(D+m),$$

$$(23) \quad p(D)y^m = y^m p(D-m).$$

The m -th powers, $m \geq 1$, of x and y satisfy

$$(24) \quad x^m y^m = \prod_{i=0}^{m-1} (D - i),$$

$$(25) \quad y^m x^m = \prod_{i=1}^m (D + i).$$

This can be seen using induction:

$$x^{m+1} y^{m+1} = x^m D y^m \stackrel{(23)}{=} x^m y^m (D - m)$$

and

$$y^{m+1} x^{m+1} = y^m (D + 1) x^m = y^m D x^m + y^m x^m \stackrel{(22)}{=} y^m x^m (D + (m + 1)).$$

Definition 3. The weighted degree of a monomial $x^m y^n \in \mathbf{F}\langle x, y \rangle$ is the integer $m - n$. A polynomial in $\mathbf{F}\langle x, y \rangle$ is said to be homogeneous of weighted degree j if all its constituting monomials have weighted degree j .

Left multiplication by xy is *homogeneity preserving*, i.e., for all $j \in \mathbf{Z}$ it maps the set of homogeneous monomials of weighted degree j into itself. To prove this, first consider a monomial $x^m y^n$ of dweighted egree $j = m - n$. Then,

$$(xy)x^m y^n \stackrel{(20)}{=} (x^m(xy) + mx^m)y^n = x^m(xy)y^n + mx^m y^n = x^{m+1}y^{n+1} + mx^m y^n,$$

and we see that weighted degree of homogeneity is indeed preserved. The general case follows immediately. Through (24) we conclude that left multiplication $x^k y^k$ is homogeneity preserving as well, for all non-negative integers k . We claim that left multiplication by $x^i y^j$ is homogeneity preserving only if $i = j$. To see this note that

$$yx^i y^j = x^i y^{i+j} + ix^{i-1} y^j$$

from which we can deduce that

$$x^m y^M x^n y^N = x^{n+m} y^{N+M} + \text{lower order terms.}$$

From which the claim follows.

Finally, a polynomial is homogeneity preserving if and only if all of its constituting monomials have this property. If this were not to be the case we could look at the highest-order non-homogeneous term and note from above $x^m y^M x^n y^N$ would give terms of a lower order in the polynomial expansion which cannot cancel as they have different powers of x or y .

It follows from these observations that

$$(26) \quad F_0 := \left\{ \sum_{n=0}^N c_n x^n y^n \mid N \in \mathbf{N}, c_1, \dots, c_N \in \mathbf{F} \right\}$$

is precisely the set of *homogeneity preserving polynomials* in $\mathbf{F}\langle x, y \rangle$.

Now everything is in place to give the proof of lemma 2.

Proof of lemma 2. As $(xy)^k$ is homogeneity preserving, we infer that there are coefficients a_i^k in $\mathbf{F}$ such that

$$(27) \quad (xy)^k = \sum_{i=0}^k a_i^k x^i y^i.$$

We will apply x^j to the right on both sides of (27) and derive an expression for the a_i^k . First note that (22) gives

$$(xy)^k x^j = x^j (xy + j)^k,$$

and (24) together with (22) gives

$$x^i y^i x^j \stackrel{(24)}{=} \prod_{\ell=0}^{i-1} (xy - \ell) x^j \stackrel{(22)}{=} x^j \prod_{\ell=0}^{i-1} (xy - \ell + j).$$

Hence, to find the coefficients a_i^k it is sufficient to consider

$$(xy + j)^k = \sum_{i=0}^k a_i^k \prod_{\ell=0}^{i-1} (xy - \ell + j).$$

Comparing the constant terms on both the left-hand side and right-hand side, we find

$$(28) \quad j^k = \sum_{i=0}^k a_i^k \prod_{\ell=0}^{i-1} (j - \ell) = \sum_{i=0}^k a_i^k (j)_i,$$

where $(j)_i$ is the falling factorial as in (16). Comparing (28) with the generating function of the Stirling numbers of the second kind $\{k_i\}$ as given in (18), we see that $a_i^k = \{k_i\}$. This concludes the proof of lemma 2. $\square$

5. The integral kernel of $L^N e^{tL}$

As mentioned before, as a first step we would like to apply D_N to the generating function $g(x, t) := e^{-2tx+t^2} = e^{-(x-t)^2+x^2}$ for the Hermite polynomials (6). We first compute the action of ∂_t^N on the generating function.

Lemma 3. *We have*

$$(29) \quad \partial_t^N e^{-(x-t)^2+x^2} = e^{-(x-t)^2+x^2} H_N(x-t).$$

Proof. We first note that,

$$\partial_t e^{-(x-t)^2} = 2(x-t)e^{-(x-t)^2} = -\partial_x e^{-(x-t)^2}.$$

Using this we get

$$\begin{aligned} \partial_t^N e^{-(x-t)^2+x^2} &= e^{x^2} \partial_t^N e^{-(x-t)^2} \\ &= e^{x^2} \partial_t^{N-1} \partial_t e^{-(x-t)^2} \\ &= -e^{x^2} \partial_t^{N-1} \partial_x e^{-(x-t)^2} \\ &= (-)^2 e^{x^2} \partial_t^{N-2} \partial_x^2 e^{-(x-t)^2} \\ &= \dots \\ &= (-1)^N e^{x^2} \partial_x^N e^{-(x-t)^2}. \end{aligned}$$

By a change of variables,

$$\partial_t^N e^{-(x-t)^2+x^2} = (-1)^N e^{-(x-t)^2+x^2} e^{(x-t)^2} \partial_y^N e^{-y^2} \Big|_{y=x-t}.$$

Hence, by (5),

$$\partial_t^N e^{-(x-t)^2+x^2} = e^{-(x-t)^2+x^2} H_N(x-t).$$

$\square$

Lemma 4. *For all $x \in \mathbf{R}$ and $t > 0$ we have*

$$(30) \quad L^N e^{-(x-t)^2+x^2} = (-1)^N e^{-(x-t)^2+x^2} \sum_{n=0}^N \left\{ \begin{matrix} N \\ n \end{matrix} \right\} t^n H_n(x-t).$$

Proof. This is now easy to prove. Recalling that $L = -t\partial_t$ and using (15), we get

$$\begin{aligned} L^N e^{-(x-t)^2+x^2} &= D_N e^{-(x-t)^2+x^2} \\ &= (-1)^N \sum_{n=0}^N \left\{ \begin{matrix} N \\ n \end{matrix} \right\} t^n \partial_t^n e^{-(x-t)^2+x^2} \\ &= (-1)^N e^{-(x-t)^2+x^2} \sum_{n=0}^N \left\{ \begin{matrix} N \\ n \end{matrix} \right\} t^n H_n(x-t). \end{aligned}$$

□

Our next theorem is the main result of this paper and provides an explicit expression for the integral kernel of $L^N e^{tL}$.

Theorem 1. *Let L be the Ornstein-Uhlenbeck operator on $L^2(\mathbf{R}^d, d\gamma)$, let $t > 0$, and let $N \geq 0$ be an integer. The integral kernel M_t^N of $L^N e^{tL}$ is given by*

$$(31) \quad \begin{aligned} M_t^N(x, y) &= M_t(x, y) \sum_{|n|=N} \binom{N}{n_1, \dots, n_d} \prod_{i=1}^d \sum_{m_i=0}^{n_i} \sum_{\ell_i=0}^{m_i} 2^{-m_i} \left\{ \begin{matrix} n_i \\ m_i \end{matrix} \right\} \binom{m_i}{\ell_i} \\ &\quad \times \left(-\frac{e^{-t}}{\sqrt{1-e^{-2t}}} \right)^{2m_i-\ell_i} H_{\ell_i}(x_i) H_{2m_i-\ell_i} \left(\frac{x_i e^{-t} - y_i}{\sqrt{1-e^{-2t}}} \right). \end{aligned}$$

Proof. We first prove the result for $d = 1$. Pulling L^N through the integral expression (4) for e^{tL} involving the Mehler kernel, we must find a suitable expression for the kernel $M_t^N(\cdot, y) = L^N M_t(\cdot, y)$. Using (7) and the normalization of H_m in (11) we get

$$\begin{aligned} M_t^N(x, y) &= L^N \sum_{m=0}^{\infty} \frac{e^{-tm}}{m!} \frac{1}{2^m} H_m(x) H_m(y) \\ &\stackrel{(8)}{=} L^N \sum_{m=0}^{\infty} \frac{e^{-tm}}{m!} \frac{1}{2^m} H_m(x) \frac{(-2i)^m}{\sqrt{\pi}} e^{y^2} \int_{-\infty}^{\infty} e^{-\xi^2} \xi^m e^{2iy\xi} d\xi \\ &= L^N \frac{e^{y^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-\xi^2} e^{2iy\xi} \sum_{m=0}^{\infty} \frac{1}{m!} H_m(x) (-i\xi e^{-t})^m d\xi \\ &\stackrel{(6)}{=} L^N \frac{e^{y^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-\xi^2} e^{2iy\xi} e^{-(x+i\xi e^{-t})^2+x^2} d\xi. \end{aligned}$$

The operator L^N is applied with respect to x here, so by lemma 4 we get

$$\begin{aligned} M_t^N(x, y) &= \frac{e^{y^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-\xi^2} e^{2iy\xi} L^N e^{-(x+i\xi e^{-t})^2+x^2} d\xi \\ &\stackrel{(30)}{=} (-1)^N \frac{e^{x^2+y^2}}{\sqrt{\pi}} \sum_{m=0}^N \left\{ \begin{matrix} N \\ m \end{matrix} \right\} \int_{-\infty}^{\infty} e^{2iy\xi} e^{-(x+i\xi e^{-t})^2} (i\xi e^{-t})^m H_m(y + i\xi e^{-t}) e^{-\xi^2} d\xi, \end{aligned}$$

where in last line we have used the analytic continuation of the algebraic identity (30). Similarly we can expand $H_m(y + i\xi e^{-t})$ using (9). This gives

$$H_m(y + i\xi e^{-t}) = \sum_{\ell=0}^m \binom{m}{\ell} H_\ell(y) (2i\xi e^{-t})^{m-\ell},$$

so that M_t^N can be written as

$$(32) \quad (-1)^N \frac{e^{x^2+y^2}}{\sqrt{\pi}} \sum_{m=0}^N \sum_{\ell=0}^m \left\{ \begin{matrix} N \\ m \end{matrix} \right\} \binom{m}{\ell} H_\ell(y) 2^{m-\ell} \int_{-\infty}^{\infty} e^{2iy\xi - \xi^2} e^{-(x+i\xi e^{-t})^2} (i\xi e^{-t})^{2m-\ell} d\xi.$$

If we set $M = 2m - \ell$, this reduces our task to computing the integral

$$(33) \quad \begin{aligned} & \frac{e^{x^2+y^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{2iy\xi - \xi^2} e^{-(x+i\xi e^{-t})^2} (i\xi e^{-t})^M d\xi \\ &= \frac{e^{y^2}}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{2i(xe^{-t}-y)\xi} e^{-(1-e^{-2t})\xi^2} (i\xi e^{-t})^M d\xi. \end{aligned}$$

To make the computation less convolved, let us set

$$\alpha_t := \sqrt{1 - e^{-2t}}, \text{ and, } \beta_t(x, y) := \frac{xe^{-t} - y}{\sqrt{1 - e^{-2t}}}.$$

This allows us to write the exponential in the integral (33) as

$$e^{2i(xe^{-t}-y)\xi} e^{-(1-e^{-2t})\xi^2} = e^{2i\alpha_t\beta_t(x,y)\xi} e^{-\alpha_t^2\xi^2}.$$

This reduces the problem, after the substitution $\alpha_t\xi \rightarrow \xi$, to computing the integral

$$\frac{e^{y^2}}{\sqrt{\pi}} \frac{i^M e^{-Mt}}{\alpha_t^{M+1}} \int_{-\infty}^{\infty} e^{2i\beta_t(x,y)\xi} e^{-\xi^2} \xi^M d\xi.$$

The final integral is an integral expression for the Hermite polynomials (8), so

$$\begin{aligned} & \frac{e^{y^2}}{\sqrt{\pi}} \frac{i^M e^{-Mt}}{\alpha_t^{M+1}} \int_{-\infty}^{\infty} e^{2i\beta_t(x,y)\xi} e^{-\xi^2} \xi^M d\xi \\ & \stackrel{(8)}{=} e^{y^2 - \beta_t(x,y)^2} \frac{1}{\alpha_t^{M+1}} \frac{(-1)^M e^{-Mt}}{2^M} H_M(\beta_t(x, y)). \end{aligned}$$

Next we note that $\exp(y^2 - \beta_t(x, y)^2) \alpha_t^{-1} = M_t$, the Mehler kernel from (4). Hence,

$$(34) \quad \begin{aligned} M_t^N(x, y) &= M_t(x, y) \sum_{m=0}^N \sum_{\ell=0}^m \binom{m}{\ell} \left\{ \begin{matrix} N \\ m \end{matrix} \right\} \left(-\frac{e^{-t}}{\sqrt{1 - e^{-2t}}} \right)^{2m-\ell} 2^{-m} \\ &\quad \times H_\ell(x) H_{2m-\ell} \left(\frac{xe^{-t} - y}{\sqrt{1 - e^{-2t}}} \right). \end{aligned}$$

Applying (13) we get the result in d dimensions. $\square$

6. An application

As an application of our main result, in this section we give an alternative proof of the bounds on the kernels K and $\tilde{K}$ of [11] (see the definition below), making the dependence on the parameters more explicit. These kernels play a central role in the study of the Hardy space $H^1(\mathbf{R}^d, d\gamma)$ in [11], where the standard Calderón reproducing formula is replaced by

$$(35) \quad u = C \int_0^\infty (t^2 L)^{N+1} e^{\frac{t^2}{\alpha} L} u \frac{dt}{t} + \int_{\mathbf{R}^d} u d\gamma,$$

where C is a suitable constant only depending on N and α (this can be seen by letting u be a Hermite polynomial). The kernels K and $\tilde{K}$ then occur in several decompositions, and the estimates below allow them to be related to classical results about the Mehler kernel.

Definition 4. *We define the kernels K and $\tilde{K}$ by*

$$\begin{aligned} \int_{\mathbf{R}^d} K_{t^2, N, \alpha}(x, y) u(y) d\gamma(y) &= (t^2 L)^N e^{\frac{t^2}{\alpha} L} u(x), \\ \int_{\mathbf{R}^d} \tilde{K}_{t^2, N, \alpha, j}(x, y) u(y) d\gamma(y) &= (t^2 L)^N e^{\frac{t^2}{\alpha} L} t \partial_{x_j}^* u(x). \end{aligned}$$

Note that the operators on the right-hand sides are indeed given by integral kernels: the first is a scaled version of the operator we have already been studying, and a duality argument implies that the second is given by the integral kernel

$$\tilde{K}_{t^2, N, \alpha, j}(x, y) = t \partial_{x_j} K_{t^2, N, \alpha}(x, y).$$

Thus, both kernels are given as appropriate derivatives of the Mehler kernel.

We begin with a technical lemma which is a rephrased version of [11, Lemma 3.4]. One should take note that we define the kernels with respect to the Gaussian measure whereas, [11] defines these with respect to the Lebesgue measure.

Lemma 5. *For all $\alpha > 1$ and all $t > 0$ and x, y in $\mathbf{R}^d$ we have*

$$(36) \quad \frac{|e^{-\frac{t}{\alpha} x} - y|^2}{1 - e^{-2\frac{t}{\alpha}}} \geq \frac{\alpha}{2} e^{-2t} \frac{|e^{-t} x - y|^2}{1 - e^{-2t}} - \frac{t^2 \min(|x|^2, |y|^2)}{1 - e^{-2\frac{t}{\alpha}}}.$$

Additionally, we have

$$(37) \quad \alpha e^{-2t} \leq \frac{1 - e^{-2t}}{1 - e^{-2\frac{t}{\alpha}}} \leq \alpha.$$

Theorem 2. *Let N be a positive integer, $0 < t < T$. The for all large enough $\alpha > 1$ we have*

(1) *If $t|x| \leq C$, then*

$$|K_{t^2, \alpha, N}(x, y)| \lesssim_{T, N} \alpha \exp\left(\frac{\alpha}{2} C^2\right) M_{t^2}(x, y) \exp\left(-\frac{\alpha}{8e^{2T}} \frac{|e^{-t^2} x - y|^2}{1 - e^{-2t^2}}\right).$$

(2) *If $t|x| \leq C$, then*

$$|\tilde{K}_{t^2, \alpha, N, j}(x, y)| \lesssim_{T, N} \alpha \exp\left(\frac{\alpha}{2} C^2\right) M_{t^2}(x, y) \exp\left(-\frac{\alpha}{8e^{2T}} \frac{|e^{-t^2} x - y|^2}{1 - e^{-2t^2}}\right).$$

Proof. For $K_{t^2, \alpha, N}$, we use Theorem 1 to obtain, after taking absolute values,

$$\begin{aligned} |K_{t^2, \alpha, N}(x, y)| &\leq M_{\frac{t^2}{\alpha}}(x, y) \sum_{|k|=N} \binom{N}{n_1, \dots, n_d} \prod_{i=1}^d t^{2k_i} \sum_{\ell_i=0}^{n_i} \sum_{m_i=0}^{m_i} 2^{-m_i} \binom{m_i}{\ell_i} \left\{ \begin{matrix} n_i \\ m_i \end{matrix} \right\} \\ &\quad \times \left(\frac{e^{-\frac{t^2}{\alpha}}}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right)^{2m_i - \ell_i} |H_{\ell_i}(x_i)| \left| H_{2m_i - \ell_i} \left(\frac{x_i e^{-\frac{t^2}{\alpha}} - y_i}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right) \right|. \end{aligned}$$

Recalling that $\ell_1 + \dots + \ell_d \leq N$, using the assumptions $t \leq T$ and $t|x| \leq C$ we can bound $t^{2k_i} |H_{\ell_i}(x)|$ by considering the highest order term to obtain

$$t^{2k_i} |H_{\ell_i}(x)| \lesssim_{C, N, T} 1.$$

Using (37) we proceed by looking at

$$\begin{aligned} M_{\frac{t^2}{\alpha}}(x, y) &= M_{t^2}(x, y) \left(\frac{1 - e^{-2t^2}}{1 - e^{-2\frac{t^2}{\alpha}}} \right)^{1/2} \exp \left(\frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(-\frac{|e^{-\frac{t^2}{\alpha}}x - y|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \\ &\leq \alpha M_{t^2}(x, y) \exp \left(\frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \left[\exp \left(-\frac{1}{2} \frac{|e^{-\frac{t^2}{\alpha}}x - y|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \right]^2. \end{aligned}$$

We can now bound the final Hermite polynomial in the expression of the kernel. Setting $M_i = 2m_i - \ell_i$ we get

$$\begin{aligned} \left(\frac{e^{-\frac{t^2}{\alpha}}}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right)^{M_i} \left| H_{M_i} \left(\frac{e^{-\frac{t^2}{\alpha}}x_i - y_i}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right) \right| &\lesssim_N \left(\frac{e^{-\frac{t^2}{\alpha}}}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right)^{M_i} \left(\frac{|e^{-\frac{t^2}{\alpha}}x_i - y_i|}{\sqrt{1 - e^{-2\frac{t^2}{\alpha}}}} \right)^{M_i} \\ &\leq \left(\frac{|e^{-\frac{t^2}{\alpha}}x_i - y_i|}{1 - e^{-2\frac{t^2}{\alpha}}} \right)^{M_i}. \end{aligned}$$

Also,

$$\left(\frac{|e^{-\frac{t^2}{\alpha}}x_i - y_i|}{1 - e^{-2\frac{t^2}{\alpha}}} \right)^{M_i} \exp \left(-\frac{1}{2} \frac{|e^{-\frac{t^2}{\alpha}}x_i - y_i|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \lesssim 1.$$

Putting these estimates together, using Lemma 5, and taking $\alpha > 1$ so large that

$$1 - \frac{\alpha}{4e^{2T}} \leq -\frac{\alpha}{8e^{2T}} \quad \text{and} \quad 1 - e^{-2\frac{t^2}{\alpha}} \geq \frac{t^2}{\alpha},$$

we obtain

$$\begin{aligned} |K_{t^2, \alpha, N}(x, y)| &\lesssim_{T, N} \alpha M_{t^2}(x, y) \exp \left(\frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(-\frac{1}{2} \frac{|e^{-\frac{t^2}{\alpha}}x - y|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \\ &\leq \alpha M_{t^2}(x, y) \exp \left(\frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(-\frac{\alpha}{4e^{2T}} \frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(\frac{1}{2} \frac{t^4|x|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \\ &= \alpha M_{t^2}(x, y) \exp \left(\left(1 - \frac{\alpha}{4e^{2T}} \right) \frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(\frac{1}{2} \frac{t^4|x|^2}{1 - e^{-2\frac{t^2}{\alpha}}} \right) \\ &\leq \alpha M_{t^2}(x, y) \exp \left(-\frac{\alpha}{8e^{2T}} \frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right) \exp \left(\frac{\alpha}{2} t^2|x|^2 \right) \\ &\leq \alpha \exp \left(\frac{\alpha}{2} C^2 \right) M_{t^2}(x, y) \exp \left(-\frac{\alpha}{8e^{2T}} \frac{|e^{-t^2}x - y|^2}{1 - e^{-2t^2}} \right), \end{aligned}$$

using $t|x| \leq C$ in the last step.

For the bound on $\tilde{K}$ we consider

$$\begin{aligned} t\partial_{x_i} \left[H_{\ell_i}(x_i) H_{m_i} \left(\frac{x_i e^{-t} - y_i}{\sqrt{1 - e^{-2t}}} \right) \right] &= t H_{m_i} \left(\frac{x_i e^{-t} - y_i}{\sqrt{1 - e^{-2t}}} \right) \partial_{x_i} H_{\ell_i}(x_i) \\ &\quad + H_{\ell_i}(x_i) t \partial_{x_i} H_{m_i} \left(\frac{x_i e^{-t} - y_i}{\sqrt{1 - e^{-2t}}} \right). \end{aligned}$$

So, as the first term on the right-hand side just decreases in degree we look at

$$t\partial_{x_i} \left(\frac{x_i e^{-t} - y_i}{\sqrt{1 - e^{-2t}}} \right)^{m_i} = m_i \left(\frac{x_i e^{-t} - y_i}{\sqrt{1 - e^{-2t}}} \right)^{m_i - 1} t \frac{e^{-t}}{\sqrt{1 - e^{-2t}}}$$

The last term is bounded as $t \downarrow 0$, and the rest of the proof is as before. $\square$

Acknowledgments. This work was partially supported by NWO-VICI grant 639.033.604 of the Netherlands Organisation for Scientific Research (NWO).

The author wishes to thank Alex Amenta and Mikko Kemppainen for inspiring discussions.

References

- [1] *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, Release 1.0.13 of 2016-09-16. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller and B. V. Saunders, eds.
- [2] E. Harboure, J.L. Torrea, and B. Viviani. On the search for weighted inequalities for operators related to the Ornstein-Uhlenbeck semigroup. *Mathematische Annalen*, 318(2):341–353, October 2000.
- [3] M. Kemppainen. An L^1 -estimate for certain spectral multipliers associated with the ornstein-uhlenbeck operator (preprint). 2015.
- [4] J. Maas, J. van Neerven, and P. Portal. Conical square functions and non-tangential maximal functions with respect to the gaussian measure. *Publicacions Matemàtiques*, 55(2):313–341, March 2011.
- [5] J. Maas, J. van Neerven, and P. Portal. Whitney coverings and the tent spaces $T_{1,q}(\gamma)$ for the Gaussian measure. *Arkiv for Matematik*, 50(2):379–395, April 2012.
- [6] G. Mauceri and S. Meda. BMO and H^1 for the ornstein-uhlenbeck operator. *J. Funct. Anal.*, 252(1):278–313, 2007.
- [7] G. Mauceri, S. Meda, and P. Sjögren. Endpoint estimates for first-order Riesz transforms associated to the ornstein-uhlenbeck operator. *Rev. Mat. Iberoam.*, 28(1):77–91, 2012.
- [8] B. Muckenhoupt. Hermite conjugate expansions. *Trans. Amer. Math. Soc.*, 139:243–260, 1969.
- [9] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST Handbook of Mathematical Functions*. Cambridge University Press, New York, NY, 2010. Print companion to [1].
- [10] E. Pineda and W. Urbina R. Non tangential convergence for the Ornstein-Uhlenbeck semigroup. *Divulgaciones Matemáticas*, 16(2):107–124, 2008.
- [11] P. Portal. Maximal and quadratic gaussian Hardy spaces. *Revista Matemática Iberoamericana*, 30:79–108, March 2014.
- [12] P. Sjögren. Operators associated with the Hermite semigroup—a survey. *J. Fourier Anal. Appl.*, 3:813–823, 1997.
- [13] J. Teuwen. A note on gaussian maximal functions. *Indagationes Mathematicae*, 26(1):106 – 112, 2015.
- [14] S. Thangavelu. *Lectures on Hermite and Laguerre expansions*, volume 42 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1993. With a preface by Robert S. Strichartz.
- [15] J. H. van Lint and R. M. Wilson. *A course in combinatorics*. Cambridge University Press, Cambridge, second edition, 2001.

Delft Institute of Applied Mathematics, Delft University of Technology, P.O. Box 5031, 2600
GA Delft, The Netherlands, j.j.b.teuwen@tudelft.nl